

On analytic properties of entropy rate

Alexander Schönhuth

Pacific Institute for the Mathematical Sciences
 School of Computing Science
 Simon Fraser University
 8888 University Drive
 Burnaby, BC, V5A 1S6, Canada
 schoenhuth@cs.sfu.ca

Abstract. Entropy rate is a real valued functional on the space of discrete random sources for which it exists. However, it lacks existence proofs and/or closed formulas even for classes of random sources which have intuitive parameterizations. A good way to overcome this problem is to examine its analytic properties relative to some reasonable topology. A canonical choice of a topology is that of the norm of total variation as it immediately arises with the idea of a discrete random source as a probability measure on sequence space. It is shown that both upper and lower entropy rate, hence entropy rate itself if it exists, are Lipschitzian relative to this topology, which, by well known facts, is close to differentiability. An application of this theorem leads to a simple and elementary proof of the existence of entropy rate of random sources with finite evolution dimension. This class of sources encompasses arbitrary hidden Markov sources and quantum random walks.

Keywords. Analytic properties, discrete random source, entropy rate, evolution dimension, hidden Markov source, quantum random walk

1 Introduction

Entropy rate is a key quantity in information theory as it is equal to the average amount of information per symbol of discrete-time, discrete-valued stochastic processes (usually referred to as *discrete random sources* in the following). Therefore, it is natural to ask how entropy rate behaves if knowledge of discrete random sources is subject to uncertainties which, for example, may be inherent to inference processes and/or originate from noisy channels. However, closed formulas for entropy rate exist only for rare examples of classes of discrete random sources. For instance, already hidden Markov sources (HMSs) seem to defy a convenient formula although there is one for the special case of Markov sources. Therefore, in this case, recent efforts focused on the direct investigation of analytic properties of entropy rate like smoothness or even analyticity [20,21], [31,30], [25], [18].

The purpose of this paper is to contribute to the issue of analytic properties of entropy rate in a more general fashion. Namely, we study the behavior of entropy rate relative to the topology induced by the norm of total variation. This topology is one of

the natural choices and it is ubiquitous in both theoretical and practical work. We show that entropy rate is Lipschitzian on the whole space of discrete random sources which is, due to an elementary theorem of Rademacher, close to differentiability.

We will use this result to give an elementary proof of the existence of entropy rate for sources with finite evolution dimension [6] which contain the classes of arbitrary HMSs [24] and quantum random walks (QRWs)[1], [5].

The paper is organized as follows. We will identify discrete random sources with probability measures acting on the measurable space of symbol sequences equipped with the σ -algebra generated by the cylinder sets of sequences. Therefore, in section 2, we will briefly compile the theory's standard arguments. In section 3 we prove that entropy rate is Lipschitz continuous relative to the topology induced by the norm of total variation which is the main contribution of this paper. In section 4 we demonstrate how to exploit this result for an elementary proof of existence of random sources with finite evolution dimension which include HMSs and QRWs as special cases. In section 5 we will describe the proof's intuition thereby commenting on open problems such as other choices of topology and/or stricter choices of analytic properties.

2 Random sources and entropy rate

As usual, $\Sigma^* = \cup_{t \geq 0} \Sigma^t$ is the set of all words (strings of finite length) over the finite alphabet Σ together with the concatenation operation

$$v \in \Sigma^t, w \in \Sigma^s \Rightarrow vw \in \Sigma^{t+s}. \quad (1)$$

Throughout this paper $\Omega = \Sigma^{\mathbb{N}} = \bigotimes_{t=0}^{\infty} \Sigma$ is the set of sequences over Σ and $\mathcal{B}$ is the σ -algebra generated by the cylinder sets. Cylinder sets B are identified with sets of words $A_B \subset \Sigma^t$ such that B is the set of sequences which start with the words in A_B . In general, the cardinality of a set A is denoted by $|A|$.

We view stochastic processes $(X_t)_{t \in \mathbb{N}}$ with values in Σ as probability measures P_X on the measurable space $(\Omega, \mathcal{B})$ and vice versa via the relationship ($v = v_0 \dots v_{t-1} \in \Sigma^t$ corresponds to the cylinder set of sequences having v as prefix)

$$P_X(v) = P(\{X_0 = v_0, X_1 = v_1, \dots, X_{t-1} = v_{t-1}\}), \quad (2)$$

where the term on the right hand side is the probability that the random source emits the symbols $v_0, \dots, v_{t-1}$ at periods $0, \dots, t-1$. Note that a stochastic process (X_t) is uniquely determined by the values $P_X(v)$ for all $v \in \Sigma^*$ as the cylinder sets corresponding to words v generate $\mathcal{B}$.

Although being a canonical choice of norm (see appendix A for a short review of the related theory and corresponding definitions), computation of the norm of total variation would not be easy for the measurable space under consideration by means of its original definition alone. The following lemma shows a concrete way to get a grip of the corresponding topology. Exact definition and basic properties of the norm of total variation have been deferred to appendix A.

Lemma 1. *The topology induced by the norm of total variation is that of the metric*

$$d_{TV}(P_X, P_Y) = \sup_{t \in \mathbb{N}} \sum_{v \in \Sigma^t} |P_X(v) - P_Y(v)| = \lim_{t \rightarrow \infty} \sum_{v \in \Sigma^t} |P_X(v) - P_Y(v)|. \quad (3)$$

where P_X, P_Y are probability measures associated to discrete random sources $(X_t), (Y_t)$.

Proof. See sec. A.2 of the appendix for the predominantly measure theoretical arguments. $\diamond$

2.1 Entropy Rate

In the following, we will refer to the quantities

$$\overline{H}(X) := \overline{H}(P_X) := \limsup_{t \rightarrow \infty} H^t(P_X) \quad (4)$$

$$\text{resp. } \underline{H}(X) := \underline{H}(P_X) := \liminf_{t \rightarrow \infty} H^t(P_X) \quad (5)$$

as *upper entropy rate* resp. *lower entropy rate* of a random source (X_t) with associated probability measure P_X , where, using the language introduced above,

$$H^t(P_X) := -\frac{1}{t} \sum_{v \in \Sigma^t} P_X(v) \log P_X(v) \quad (6)$$

is the entropy of the distribution over the words of length t induced by the random source, divided by t . *Entropy rate* of a random source (X_t) with associated probability measure P_X is denoted by

$$H(X) := H(P_X) := \lim_{t \rightarrow \infty} H^t(P_X). \quad (7)$$

The existence of the limit of the $H^t(P_X)$ is also referred to as the *existence of entropy rate* where, obviously, a necessary and sufficient condition for entropy rate to exist is

$$\overline{H}(X) = \underline{H}(X) \quad (= H(X)). \quad (8)$$

Throughout this paper, $\Delta^{n-1} = \{x = (x_1, \dots, x_n) \in \mathbb{R}^n \mid x_i \geq 0, \sum_i x_i = 1\}$ is the usual regular $n - 1$ -dimensional simplex in $\mathbb{R}^n$ and, for technical convenience, $\log$ is the natural logarithm. Note that, as it is more common to use the logarithm to the base 2, switching bases does not affect any analytic property of entropy rate.

3 Analytic properties of entropy rate

Our main result is the following theorem, which states that entropy rate is Lipschitz continuous with respect to the topology induced by the norm of total variation. In the following let $\mathcal{P}$ be the set of the probability measures associated with discrete random sources, viewed as a normed space. Elements of $\mathcal{P}$ will be denoted by P or Q . We further denote the normed subspace of discrete random sources for which entropy rate exists by $\mathcal{P}_H$.

Theorem 1 (Lipschitz continuity of entropy rate). *The real-valued functionals $\overline{H}$ and $\underline{H}$ on $\mathcal{P}$ are Lipschitzian with $Lip(\overline{H}) = Lip(\underline{H}) = \log |\Sigma|$, that is, for $P, Q \in \mathcal{P}$,*

$$|\overline{H}(P) - \overline{H}(Q)| \leq (\log |\Sigma|) d_{TV}(P, Q) \quad (9)$$

$$|\underline{H}(P) - \underline{H}(Q)| \leq (\log |\Sigma|) d_{TV}(P, Q). \quad (10)$$

Clearly, because of (8), a corollary of the theorem is that the same holds true for entropy rate itself.

Corollary 1. *Entropy rate is Lipschitzian with $Lip(H) = \log |\Sigma|$, that is,*

$$|H(P) - H(Q)| \leq (\log |\Sigma|) d_{TV}(P, Q) \quad (11)$$

where, here, $P, Q \in \mathcal{P}_H$.

We present two lemmata, which incorporate the essential ideas of the proof of the theorem. We write

$$d_{TV,t}(P, Q) := \sum_{v \in \Sigma^t} |P(v) - Q(v)|. \quad (12)$$

Lemma 1 says that $\lim_{t \rightarrow \infty} d_{TV,t}(P, Q) = d_{TV}(P, Q)$. Note that $d_{TV,t}$ is not a metric on $\mathcal{P}$.

Lemma 2. *Let $P, Q \in \mathcal{P}$ such that $d_{TV}(P, Q) \leq \frac{1}{e}$. Then it holds that*

$$|H^t(P) - H^t(Q)| \leq (\log |\Sigma| + \frac{1}{t} \log \frac{1}{d_{TV,t}(P, Q)}) \cdot d_{TV,t}(P, Q),$$

where $0 \cdot \log \infty := 0$ in case of $d_{TV,t}(P, Q) = 0$.

For the proof of this lemma we will need a technical sublemma.

Sublemma 31 *Let $h(x) := x \log(1/x)$ for $x \in]0, 1]$ and $h(0) = 0$. Then, for $x, y \in [0, 1]$,*

$$|x - y| \leq \frac{1}{e} \implies |h(x) - h(y)| \leq h(|x - y|). \quad (13)$$

Proof. Note first that $h'(x) = \log \frac{1}{x} - 1$ and $h''(x) = -\frac{1}{x}$. Hence h is concave, has a global maximum at $\frac{1}{e}$ and $h(\frac{1}{e}) = \frac{1}{e}$. Therefore $x \leq h(x) \iff x \leq \frac{1}{e} (*)$. Because of

$$\begin{aligned} |h(x) - h(y)| &= |h(x) - h(\frac{1}{e})| - |h(\frac{1}{e}) - h(y)| \\ &\leq \max\{|h(x) - h(\frac{1}{e})|, |h(\frac{1}{e}) - h(y)|\} \end{aligned} \quad (14)$$

and the fact that h is monotonically increasing on $[0, \frac{1}{e}]$ we can, without loss of generality, assume that either $x, y \geq \frac{1}{e}$ or $x, y \leq \frac{1}{e}$. Because of $|h'(x)| \leq 1$ on $[\frac{1}{e}, 1]$ and the mean value theorem, it holds that

$$\frac{1}{e} \leq x, y \leq 1 \Rightarrow |h(x) - h(y)| \leq |h'(x)||x - y| = |x - y|. \quad (15)$$

Because of (*) we obtain the claim for the case $\frac{1}{e} \leq x, y \leq 1$.

It remains the case (w.l.o.g. $x < y$) $x < y \leq \frac{1}{e}$. Here it holds that $|h(x) - h(y)| = h(y) - h(x)$. We note that the function $\log \frac{1}{t} - 1$ is positive and monotonically decreasing on $[0, \frac{1}{e}]$ (**). We obtain the claim from the calculation

$$\begin{aligned} |h(x) - h(y)| &= \int_x^y (\log \frac{1}{t} - 1) dt \stackrel{(**)}{\leq} \int_x^y (\log \frac{1}{t-x} - 1) dt \\ &\stackrel{s=t-x}{=} \int_0^{y-x} (\log \frac{1}{s} - 1) ds = \left[s \log \frac{1}{s} \right]_0^{y-x} = h(y-x). \end{aligned} \quad (16)$$

◇

Let now $\Delta_K^{n-1} := K \cdot \Delta^{n-1} = \{x = (x_1, \dots, x_n) \in \mathbb{R}^n \mid x_i \geq 0, \sum_i x_i = K\}$. In a way that is completely analogous to that of showing that entropy attains a maximum at uniform distributions we infer that, on Δ_K^{n-1} , the function $h_{K,n}(x_1, \dots, x_n) := \sum_{i=1}^n x_i \log \frac{1}{x_i}$ (a scaled version of entropy) attains a global maximum at $\bar{x} := (K/n, \dots, K/n)$ (***).

We are now able to prove lemma 2.

Proof. Obviously $H^t(P) = H^t(Q)$ in case of $d_{TV,t}(P, Q) = 0$. In case of $d_{TV,t}(P, Q) > 0$

$$\begin{aligned} &|H^t(P) - H^t(Q)| \\ &\leq \frac{1}{t} \sum_{v \in \Sigma^t} |P(v) \log \frac{1}{P(v)} - Q(v) \log \frac{1}{Q(v)}| \\ &\stackrel{Subl. 31}{\leq} \frac{1}{t} \sum_{v \in \Sigma^t} |P(v) - Q(v)| \log \frac{1}{|P(v) - Q(v)|} \\ &\stackrel{(***)}{\leq} \frac{1}{t} \sum_{v \in \Sigma^t} \frac{d_{TV,t}(P, Q)}{|\Sigma|^t} \log \frac{|\Sigma|^t}{d_{TV,t}(P, Q)} \\ &= \frac{1}{t} d_{TV,t}(P, Q) (t \log |\Sigma| + \log \frac{1}{d_{TV,t}(P, Q)}). \end{aligned} \quad (17)$$

◇

To get control of the limes superior resp. inferior involved in the definition of entropy rate we will further need the following lemma.

Lemma 3. Let (a_t) and (b_t) two non-negative real valued sequences such that

$$|a_t - b_t| \leq c_t \quad \text{and} \quad \lim_{t \rightarrow \infty} c_t = c. \quad (18)$$

Then it holds that

$$|\limsup_{t \rightarrow \infty} a_t - \limsup_{t \rightarrow \infty} b_t| \leq c \quad (19)$$

$$|\liminf_{t \rightarrow \infty} a_t - \liminf_{t \rightarrow \infty} b_t| \leq c. \quad (20)$$

Proof. We only display the proof for (19) as that of (20) can be obtained, mutatis mutandis, by analogous considerations.

W.l.o.g. assume $a := \limsup_{t \rightarrow \infty} a_t \geq \limsup_{t \rightarrow \infty} b_t =: b$. Choose a subsequence $k(t)$ such that $\lim_{t \rightarrow \infty} a_{k(t)} = a$. We obtain

$$\begin{aligned} a - b &\leq a - \limsup_{t \rightarrow \infty} b_{k(t)} = \limsup_{t \rightarrow \infty} a_{k(t)} - \limsup_{t \rightarrow \infty} b_{k(t)} \\ &\leq \limsup_{t \rightarrow \infty} |a_{k(t)} - b_{k(t)}| \leq c. \end{aligned} \quad (21)$$

◇

We are now in position to prove theorem 1.

Proof. As Lipschitz continuity is a local property, we can assume that $d_{TV}(P, Q) \leq \frac{1}{e}$. Setting $a_t := H^t(P)$ and $b_t := H^t(Q)$ we obtain by lemma 2

$$|a_t - b_t| \leq d_{TV,t}(P, Q) \left(\log |\Sigma| + \frac{1}{t} \log \frac{1}{d_{TV,t}(P, Q)} \right) =: c_t. \quad (22)$$

The definition of $d_{TV,t}$ and lemma 1 lead to

$$\begin{aligned} \lim_{t \rightarrow \infty} c_t &= \lim_{t \rightarrow \infty} d_{TV,t}(P, Q) \left(\log |\Sigma| + \frac{1}{t} \log \frac{1}{d_{TV,t}(P, Q)} \right) \\ &= d_{TV}(P, Q) \cdot \log |\Sigma|. \end{aligned} \quad (23)$$

Plugging (a_t) , (b_t) and (c_t) into lemma 3 then yields the desired result. ◇

In order to elucidate that the structure of the proof strongly depends on the choice of the norm we rephrase lemma 2 in a more general fashion, without the “soul” of an entropy. Therefore let

$$h_n(x_1, \dots, x_n) = \frac{1}{\log n} \sum_{i=1}^n x_i \log \frac{1}{x_i} \quad (24)$$

on Δ^{n-1} where $n \geq 2$ and $0 \log \infty := 0$. A more prosaic version of lemma 2 then reads

$$|h_n(x) - h_n(y)| \leq \|x - y\|_1 \cdot \left(1 + \frac{1}{\log n} \log \frac{1}{\|x - y\|_1} \right), \quad (25)$$

where $\|x\|_1 = \sum_i |x_i|$ as usual. A straightforward consequence of the lemma is

$$\forall \epsilon \in \mathbb{R}_+ \exists \delta \in \mathbb{R}_+ \forall n \geq 2 \forall x, y \in \Delta^{n-1} : \|x - y\|_1 < \delta \implies |h_n(x) - h_n(y)| < \epsilon. \quad (26)$$

After being translated back to entropies, this states that entropy rate is uniformly continuous on $\mathcal{P}$. We note that the statement of the generalized lemma need not be true relative to norms $\|\cdot\|_p$ different from $\|\cdot\|_1$. More formally:

Lemma 4. *Let $2 \leq p < \infty$ and $\|x\|_p = \sqrt[p]{\sum_i |x_i|^p}$ the usual p -norm on $\mathbb{R}^n$. Then it holds that*

$$\exists \epsilon \in \mathbb{R}_+ \forall \delta \in \mathbb{R}_+ \exists N \geq 2 \exists x, y \in \Delta^{N-1} : \|x - y\|_p < \delta, |h_N(x) - h_N(y)| > \epsilon$$

which is just the negation of (26).

For the proof we use the notation ($0 < m \leq n$)

$$x_{m,n}^* := (\underbrace{\frac{1}{m}, \dots, \frac{1}{m}}_{m \text{ times}}, 0, \dots, 0) \in \Delta^{n-1}. \quad (27)$$

Proof. Choose $\epsilon = 1/2$ and $\delta \in \mathbb{R}_+$ arbitrarily. Choose an $m \in \mathbb{N}$, such that $m > \frac{1}{\delta}$. Then find an $N_0 > m$, such that $\|x_{n,n}^*\|_2 = (\frac{1}{n})^{1/2} < \delta$ for every $n \geq N_0$. Further

$$\begin{aligned} \|x_{m,n}^* - x_{n,n}^*\|_p &\leq \|x_{n,n}^*\|_p = \left(\frac{1}{n^{p-1}}\right)^{\frac{1}{p}} = n^{-\frac{p-1}{p}} \\ &= n^{\frac{1}{p}-1} \leq n^{-\frac{1}{2}} = \left\| \left(\frac{1}{n}, \dots, \frac{1}{n}\right) \right\|_2 < \delta, \end{aligned} \quad (28)$$

but

$$|h_n(x_{m,n}^*) - h_n(x_{n,n}^*)| = \frac{1}{\log n} |\log m - \log n| \xrightarrow{n \rightarrow \infty} 1. \quad (29)$$

Therefore, we find an $N \in \mathbb{N}$ and suitable $x, y \in \Delta^{N-1}$ which support the statement of the lemma. $\diamond$

REMARK Because of lemma 4, one could intuitively be led to the assumption that entropy rate need not be continuous with respect to the norms given through the spaces $L_p(\Omega, \mathcal{B}, P)$, $p \geq 2$. However, this is not true, see [27] for respective considerations.

4 Entropy rate of sources with finite evolution dimension

In the following we will give a direct proof of the existence of entropy rate of sources with finite evolution dimension which had been introduced in [6]. See the subsequent subsection 4.3 for prevalent examples of random sources of finite evolution dimension.

As sources with finite evolution dimension are asymptotically mean stationary [6], the result can be obtained as a corollary of the theorem of Shannon-McMillan-Breiman for asymptotically mean stationary sources [9]. However, the following proof is much simpler. See subsection 4.4 for a detailed comparison of the two proofs.

4.1 Preliminaries

In the following let the *shift operator* $T : \Omega \rightarrow \Omega$ be defined by

$$T(v_0 v_1 v_2 \dots) := v_1 v_2 \dots \quad (30)$$

Obviously, T is measurable. If (X_t) is a discrete random source with associated measure P_X then

$$(P_X \circ T^{-k})(v) := \sum_{w \in \Sigma^k} P_X(wv) \quad (31)$$

gives rise to a probability measure $P \circ T^{-k}$ which is associated with the discrete random source $((X_k)_t)$ defined through

$$\begin{aligned} P_{X_k}(\{(X_k)_0 = v_0, (X_k)_1 = v_1, \dots, (X_k)_{t-1} = v_{t-1}\}) \\ := P_X(\{X_k = v_0, X_{k+1} = v_1, \dots, X_{t-1+k} = v_{t-1}\}). \end{aligned}$$

A discrete random source (X_t) is said to be of *finite evolution dimension* if the family $(P_X \circ T^{-k})_{k \geq 0}$ spans a finite-dimensional subspace in the linear space of finite, signed measures on $(\Omega, \mathcal{B})$ (see appendix A for the definition of a finite, signed measure).

In the following we will write

$$PT^{-i} := P \circ T^{-i} \quad \text{and} \quad P_n := \frac{1}{n} \sum_{i=0}^{n-1} PT^{-i} \quad (32)$$

for probability measures P associated with random sources.

Theorem 2. *If P is a discrete random source of finite evolution dimension there is a stationary discrete random source $\bar{P}$, called the stationary mean of P such that*

$$\lim_{n \rightarrow \infty} d_{TV}(P_n, \bar{P}) = 0. \quad (33)$$

Proof. The proof is centered on an elementary fact from linear algebra. As it requires some of the basic theory of finite, signed measures, we have deferred it to sec. A.3 in the appendix. Note that an alternative, slightly more complicated version of the proof has already been given in [6]. $\diamond$

4.2 Proof for the existence of entropy rate

In order to be prepared for the proof we provide a lemma whose immediate consequence is that entropy rate coincides for all $P_n, n \geq 0$.

Lemma 5. *Let P be a probability measure associated with a random source. Then it holds that*

$$\forall n \in \mathbb{N} : \quad \lim_{t \rightarrow \infty} (H^t(P) - H^t(P_n)) = 0. \quad (34)$$

Proof. A straightforward consequence of Lemma 2.3.4, [10] is that for $\alpha \in [0, 1]$ and probability measures P, Q :

$$\begin{aligned} \alpha H^t(P) + (1 - \alpha) H^t(Q) &\leq H^t(\alpha P + (1 - \alpha) Q) \\ &\leq \alpha H^t(P) + (1 - \alpha) H^t(Q) + \frac{\log 2}{t} \end{aligned}$$

Now, by induction on n ,

$$\frac{1}{n} \sum_{i=0}^{n-1} H^t(P T^{-i}) \leq H^t(P_n) \leq \frac{1}{n} \sum_{i=0}^{n-1} H^t(P T^{-i}) + \frac{n}{t} \log 2$$

and the assertion follows from lemma 7 (appendix B) which states that the $H^t(P T^{-i})$ coincide for all $i \geq 0$. $\diamond$

We establish that both upper entropy rate $\overline{H}$ and lower entropy rate $\underline{H}$ coincide for all $P_n, n \geq 0$.

Corollary 2. *Let P be the probability measure associated with a random source. Then it holds that*

$$\overline{H}(P) = \overline{H}(P_n) \quad \text{and} \quad \underline{H}(P) = \underline{H}(P_n) \quad (35)$$

for all $n \in \mathbb{N}$.

Proof. Use lemma 5 in order to apply lemma 3 to the sequences $(a_t := H^t(P)), (b_t := H^t(P_n))$ for the first equation. For the second one rephrase lemma 3 with $\liminf$ instead of $\limsup$. $\diamond$

As a consequence, we can prove the existence of entropy rate for finite-evolution-dimensional sources.

Theorem 3 (Existence of entropy rate). *Let P be a probability measure associated with a random source of finite evolution dimension. Let $\bar{P}$ be the stationary mean of P . Then it holds that*

$$\overline{H}(P) = \underline{H}(P) = \lim_{t \rightarrow \infty} H^t(P). \quad (36)$$

Therefore, entropy rate of P exists. Moreover, it is equal to the one of the stationary mean $\bar{P}$.

Proof. As the P_n converge in TV-norm to $\bar{P}$ (theorem 2) we obtain due to the continuity of $\overline{H}, \underline{H}$ (theorem 1)

$$\lim_{n \rightarrow \infty} \overline{H}(P_n) = \overline{H}(\bar{P}) \quad \text{and} \quad \lim_{n \rightarrow \infty} \underline{H}(P_n) = \underline{H}(\bar{P}). \quad (37)$$

It follows, as $\overline{H}(P_n)$ and $\underline{H}(P_n)$ are constant with respect to n (corollary 2) and $\overline{H}(\bar{P}) = \underline{H}(\bar{P})$ (as entropy rate exists for stationary sources) that $\overline{H}(P) = \underline{H}(P) = \overline{H}(\bar{P})$. $\diamond$

REMARK Theorem 2 can be generalized to general asymptotically mean stationary (AMS) sources (see [9] for the theory of AMS sources). However, the proof needs a sophisticated ergodic theorem, thereby loosing the elementary flavour [26].

4.3 Examples of discrete random sources of finite evolution dimension

In the following, we present two classes of discrete random sources that have finite evolution dimension.

Hidden Markov Sources (HMSs) *Hidden Markov sources (HMSs)* are the discrete random sources associated with hidden Markov models (HMMs) (also termed hidden Markov chains in the related literature). HMSs have been largely studied, see e.g. [24] for a comprehensive review. In the following, we will give a brief definition of HMMs.

An HMM $\mathcal{M} = (\Sigma, S, \pi, A, E)$ is specified by a finite set of output symbols Σ , a set of hidden states $S = \{1, \dots, n\}$, a transition probability matrix $A = (A_{ij})_{i,j \in S} \in \mathbb{R}^{n \times n}$, an initial probability distribution $\pi \in \mathbb{R}^n$ and an emission probability matrix $E = (E_{iv})_{i \in S, v \in \Sigma} \in \mathbb{R}^{n \times \Sigma}$. It gives rise to a discrete random source $p_{\mathcal{M}}$ with values in the finite set Σ , referred to as *hidden Markov source (HMS)* by the idea of changing hidden states according to the transition probabilities $A_{ij} = P(i \rightarrow j)$, where the first state is picked according to π , and emitting symbols from the hidden states, as specified by the emission probabilities $E_{ia} = P(a \text{ is emitted from } i)$. More formally, in accordance with (2),

$$p_{\mathcal{M}}(v = v_1 \dots v_t) = \sum_{i_1 \dots i_t \in S^t} \pi(i_1) E_{i_1 v_1} A_{i_1 i_2} E_{i_2 v_2} \cdot \dots \cdot A_{i_{t-1} i_t} E_{i_t v_t}. \quad (38)$$

In the literature, HMSs are often introduced as being induced by *finite functions of Markov chains* where emission probability distributions are replaced by a finite function $f : S \rightarrow \Sigma$ mapping hidden states to output symbols. It is straightforward to see that they give rise to complete class of HMSs as well.

It is well known that HMSs have *finite dimension* or, equivalently, have *finite degree of freedom*. See [13] for an early work on the topic and [14] for further related work. The relationship of finite dimension and finite evolution dimension has been thoroughly discussed in [6]. It holds that finite evolution dimension is a necessary condition of finite dimension, which establishes that HMSs are of finite evolution dimension. Examples for which the generalization of the existence of entropy rate of sources with finite evolution dimension apply are non-stationary HMSs. A simple example for this might be a binary-valued source (i.e. $\Sigma = \{0, 1\}$) induced by a “circular” HMM acting on three hidden states $S := \{1, 2, 3\}$ with transition resp. emission probability matrix

$$\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} \quad \text{resp.} \quad \begin{bmatrix} 0 & 1 \\ 0.5 & 0.5 \\ 1 & 0 \end{bmatrix}. \quad (39)$$

Clearly, this source is not stationary such that the simple existence proof for stationary sources does not apply. However, as an HMS, this source is of finite evolution dimension such that theorem 3 ensures the existence of its entropy rate.

See the subsequent sec. 4.4 for a comparison of available proofs of the existence of entropy rate.

REMARK Related work on analytic properties of entropy rate of HMSs is concerned with topologies referring to the parameterizations of the HMMs giving rise to the

HMSs, that is, with the natural topologies of real-valued vector spaces (e.g. [20,21,22]). For example, in the special case of binary valued i.i.d. processes, emitting values from $\{0, 1\}$, entropy rate is computed as

$$p \log\left(\frac{1}{p}\right) \quad (40)$$

where the only parameter p is the probability that the binary valued i.i.d. process emits a 1. Clearly, $p \log(1/p)$ is not Lipschitz continuous in intervals around zero $[(p \log(1/p))' = \log 1/p - 1]$. However, this does not contradict theorem 1 as convergence w.r.t. the parameterization does not imply convergence w.r.t. the norm of total variation, which we will briefly outline in the following.

As follows from elementary measure theoretical considerations, the topology induced by the norm of total variation is equivalent to that of the general version of the *metric of total variation*

$$D_{TV}(P, Q) := \sup_{B \in \mathcal{B}} |P(B) - Q(B)| \quad (41)$$

where $P, Q \in \mathcal{P}$ are two probability measures acting on the measurable space $(\Omega, \mathcal{B})$. In the case of the measurable sequence spaces under consideration here, the equivalence of the topologies of the metric and the norm of total variation can be seen by lemma 1 as it follows from straightforward elementary computations that the topologies of d_{TV} of lemma 1 and the metric of total variation D_{TV} of (41) are equivalent. As a consequence, convergence in the sense of the norm of total variation is equivalent to uniform convergence on all measurable sets, that is,

$$\lim_{n \rightarrow \infty} \|P - P_n\|_{TV} = 0 \quad \Leftrightarrow \quad \lim_{n \rightarrow \infty} \sup_{B \in \mathcal{B}} |P(B) - P_n(B)| = 0 \quad (42)$$

where $P, P_n \in \mathcal{P}$.

However, as outlined in [20], sec. VIII, convergence of probability measures induced by hidden Markov models whose parameterizations converge may not even be *strong* (see [15] for definitions and characterizations of several forms of convergence of probability measures) meaning that there might exist a set $B^* \in \mathcal{B}$ for which

$$\limsup_{n \rightarrow \infty} |P(B^*) - P_n(B^*)| > 0 \quad (43)$$

where the $P_n, n = 0, 1, \dots$ are hidden Markov models whose parameterizations converge to the parameterization of P . According to (42), this means that convergence in terms of the parameterization does not necessarily imply convergence w.r.t. the norm of total variation.

Quantum Random Walks (QRWs) Quantum random walks (QRWs) were introduced to quantum information theory in 2001 as an analogon to classical Markov sources [1]. For example, they allow to emulate Markov Chain Monte Carlo approaches on quantum computers. However, their properties are much less understood. A QRW $\mathcal{Q} = (G, U, \psi_0)$, in a very general form (see [1] for the full range of definitions), is specified

by a directed, K -regular graph $G = (V, E)$, a unitary (*evolution*) operator $U : \mathbb{C}^N \rightarrow \mathbb{C}^N$ and a *wave function* $\psi_0 \in \mathbb{C}^N$ (i.e. $\|\psi_0\| = 1$ for $\|\cdot\|$ the Euclidean norm) where $N := K \cdot |V| = |E|$. Dimensions are labeled by edges which in turn are labeled by (u, x) where $u \in V$ and $x \in X$, $|X| = K$ and $\mathbb{C}^N$ is considered to be spanned by the orthonormal basis $(e_{(u,x)})_{(u,x) \in V \times X = E}$. A QRW induces a classical random source p_Q with values in $\Sigma := V$ (i.e. the set of nodes) by the following iterative procedure. In the first step, the evolution operator is applied to the initial wave function ψ_0 , and the resulting wave function $U\psi_0$, with probability $\sum_{x \in X} |(U\psi_0)_{(u_1,x)}|^2$, is collapsed (i.e. projected and renormalized, which models a quantum mechanical measurement) to the subspace of $\mathbb{C}^N$, spanned by the vectors $e_{(u_1,x)}$, $x \in X$ that is associated with (the edges leaving from) node u_1 , thereby generating the first symbol u_1 . This procedure results in a new wave function describing the state ψ_{u_1} the QRW is in after having generated the first symbol u_1 . In order to generate a second symbol U is applied to ψ_{u_1} , and $U\psi_{u_1}$ is, with probability $\sum_{x \in X} |(U\psi_{u_1})_{(u_2,x)}|^2$, collapsed to state $\psi_{u_1 u_2}$, thereby generating the second symbol u_2 . Iterative application of this basic procedure of evolving followed by collapsing yields a sequence of symbols. See [1] for further details.

A concise formal description in terms of formula analogous to (38) of the discrete random source p_Q along with a proof of QRWs being of finite dimension has been presented in [28]. Finite evolution dimension follows from finite dimension, which, as outlined above, has been thoroughly discussed in [6].

4.4 Comparison of existence proofs of entropy rate

The result of theorem 3 for the special case of HMSs can be obtained as a combination of the Shannon-McMillan-Breiman (SMB) theorem for *asymptotically mean stationary* (AMS) sources [9] and the fact that HMSs are AMS [19] (see also [24] for a comprehensive review of theoretical results on HMSs). Therefore, the existence of entropy rate for arbitrary, stationary and non-stationary, HMSs has theoretically been known since 1981. For QWRs the result has been known since 2006, implied by combining the results of [9] and [5] in the same fashion as for HMSs. However, even for HMSs, the result seems to be rather unnoticed which might be due to both the complex nature of its proof and that the necessary combination of results has not been explicitly mentioned. The SMB theorem in this most generalized version is centered around a proof for the class of ergodic, stationary random sources [29,23,3,4] which requires involved ergodic theorems. The extension to general stationary sources [2,16,17], in an exemplary (and elegant) version, needs the sophisticated concept of the ergodic decomposition of stationary random sources [8]. The final step [9] requires again a collection of non-trivial theorems as a prerequisite.

The proof given here is substantially simpler from two main aspects. First, it is centered around the standard elementary proof of the existence of entropy rate of stationary sources. Note that, this way, we do not even need to introduce ergodicity. Second, the extension to non-stationary classes of random sources is done by results of exclusively elementary nature (theorems 2, 1).

5 Conclusion

We show that entropy rate is Lipschitzian relative to the topology of total variation in an elementary fashion. Besides from providing a comparatively simple existence proof for HMSs and QRWs, this helps getting a more general grip of entropy rate. Moreover, it brings up some interesting open questions:

- A first open question which immediately arises is whether our arguments can be strengthened to stricter analytic properties. A first clue is that the definition of entropy rate as well as theorem 1 can be consistently extended to the whole real vector space of finite, signed measures. Rademacher’s theorem [7] states that Lipschitzian functionals on finite-dimensional real vector spaces are differentiable almost everywhere w.r.t. the Lebesgue measure on the Borel-sets. This points at that entropy rate is close to being differentiable and, so far, we have not succeeded in constructing a random source at which entropy rate is not differentiable.
- The intuition behind our proof is that entropy rate cannot differ too much if sets of typical sequences of two sources overlap to a sufficiently high degree. However, it seems to be obvious that entropy rate is continuous when considering it relative to sizes of sets of typical sequences which is a more general assumption. A corresponding result would certainly be applicable to coarser topologies as, say, the weak topology. So far, it has only been known that entropy rate, as a functional on the set of stationary sources only, is upper semicontinuous [10] relative to the weak topology. We believe that theorems of the quality of theorem 1, based on the comparison of sizes of sets of typical sequences, will greatly improve such results.

A The norm of total variation

In the following, let $A \dot{\cup} B$ be the disjoint union of two sets A and B and $\mathbb{C}A$ be the complement of a set A .

A.1 Finite signed measures

A finite, signed measure on $(\Omega, \mathcal{B})$ is a σ -additive but not necessarily positive, finite set function on $\mathcal{B}$. The most important relevant properties of finite signed measures are summarized in the following theorem (see [11], ch. VI for proofs).

Theorem 4.

1. *By eventwise addition and scalar multiplication, the set of finite signed measures can be considered as a real-valued vector space.*
2. *The Jordan decomposition theorem states that for every $P \in \mathcal{P}$ there are finite measures P_+, P_- such that*

$$P = P_+ - P_- \tag{44}$$

and for all other decompositions $P = P_1 - P_2$ with measures P_1, P_2 it holds that $P_1 = P_+ + \delta, P_2 = P_- + \delta$ for another measure δ . In this sense, P_+ and P_- are unique and called positive resp. negative variation. The measure $|P| := P_+ + P_-$ is called total variation.

3. In parallel to the Jordan decomposition we have the Hahn decomposition of Ω into two disjoint events Ω_+, Ω_-

$$\Omega = \Omega_+ \dot{\cup} \Omega_- \quad (45)$$

such that $P_-(\Omega_+) = 0$ and $P_+(\Omega_-) = 0$. Ω_+, Ω_- are uniquely determined up to $|P|$ -null-sets.

4. The norm of total variation $\|\cdot\|_{TV}$ on $\mathcal{P}$ is given by

$$\|P\|_{TV} := |P|(\Omega) = P_+(\Omega) + P_-(\Omega) = P_+(\Omega_+) + P_-(\Omega_-). \quad (46)$$

Obviously $\| |P| \|_{TV} = \|P\|_{TV}$.

A.2 Proof of lemma 1

For the proof, we will identify cylinder sets $B \in \mathcal{B}$ with sets of words $A_B \in \Sigma^t$ as usual (B is the set of sequences which are the continuations of the words in A_B). In our notation, we correspondingly obtain

$$P(B) = \sum_{v \in A_B} P_+(v) - P_-(v) \quad (47)$$

for a signed measure P with Jordan decomposition $P = P_+ - P_-$. We will further make use of the approximation theorem (see Halmos [11], p. 56, Th. D) which tells that, given a measure P , an event $B \in \mathcal{B}$ and $\epsilon \in \mathbb{R}_+$, we find a cylinder set F such that

$$P(B \triangle F) < \epsilon, \quad (48)$$

where $B \triangle F = (B \setminus F) \cup (F \setminus B)$ is the symmetric set difference. A straightforward consequence of this is that $|P(B) - P(F)| < \epsilon$.

Proof. It suffices to show

$$\|P\|_{TV} = \sup_{t \in \mathbb{N}} \sum_{v \in \Sigma^t} |P(v)| = \lim_{t \rightarrow \infty} \sum_{v \in \Sigma^t} |P(v)|. \quad (49)$$

for an arbitrary finite, signed measure P .

The second equation of (3) now follows immediately from

$$\sum_{v \in \Sigma^t} |P(v)| = \sum_{v \in \Sigma^t} \underbrace{\left| \sum_{a \in \Sigma} P(va) \right|}_{=|P(v)|} \leq \sum_{v \in \Sigma^t} \sum_{a \in \Sigma} |P(va)| = \sum_{v \in \Sigma^{t+1}} |P(v)| \quad (50)$$

which shows that $(\sum_{v \in \Sigma^t} |P(v)|)_{t \in \mathbb{N}}$ is a monotonically increasing sequence. It remains to show that it converges to $\|P\|_{TV}$. This translates to demonstrate that, given $\epsilon \in \mathbb{R}_+$, there is $T_0 \in \mathbb{N}$ with

$$\sum_{v \in \Sigma^{T_0}} |P(v)| > \|P\|_{TV} - \epsilon. \quad (51)$$

Therefore let P_+, P_- be the Jordan decomposition of P and, correspondingly, $\Omega = \Omega_+ \dot{\cup} \Omega_-$ be the Hahn decomposition. By an application of the approximation theorem (see above) we find $T_0 \in \mathbb{N}$ and a cylinder set corresponding to $A \subset \Sigma^{T_0}$ with

$$|P|(\Omega_+ \triangle A) < \frac{\epsilon}{4} \quad (52)$$

a straightforward ($|P| = P_+ + P_-$) consequence of which is that both

$$P_+(\Omega_+ \triangle A) < \frac{\epsilon}{4} \quad \text{and} \quad P_-(\Omega_+ \triangle A) < \frac{\epsilon}{4} \quad (53)$$

Now note that the obvious $\mathbb{C}A \triangle \mathbb{C}B = A \triangle B$ in combination with $\Omega_- = \mathbb{C}\Omega_+$ and (53) yields

$$P_-(\Omega_- \triangle \mathbb{C}A) = P(\Omega_+ \triangle A) < \frac{\epsilon}{4}. \quad (54)$$

(53) and (54) then yield the inequalities

$$P_+(\mathbb{C}A) \stackrel{P_+(\Omega_-)=0}{=} P_+(\Omega_+ \setminus A) \leq P_+(\Omega_+ \triangle A) < \frac{\epsilon}{4} \quad (55)$$

and

$$P_-(A) \stackrel{P_-(\Omega_+)=0}{=} P_-(\Omega_- \setminus \mathbb{C}A) \leq P_-(\Omega_- \triangle \mathbb{C}A) < \frac{\epsilon}{4}. \quad (56)$$

Moreover, it is straightforward from (53) and (54) that

$$P_+(A) > P_+(\Omega_+) - \frac{\epsilon}{4} \quad \text{and} \quad P_-(\mathbb{C}A) > P_-(\Omega_-) - \frac{\epsilon}{4}. \quad (57)$$

We finally compute

$$\begin{aligned} \sum_{v \in \Sigma^{T_0}} |P(v)| &= \sum_{v \in A} |P(v)| + \sum_{v \in \mathbb{C}A} |P(v)| \\ &\geq |P(A)| + |P(\mathbb{C}A)| \\ &\geq P_+(A) - P_-(A) + P_-(\mathbb{C}A) - P_+(\mathbb{C}A) \\ &\stackrel{(55), (56)}{\geq} (P_+(\Omega_+) - \frac{\epsilon}{4}) - \frac{\epsilon}{4} + (P_-(\Omega_-) - \frac{\epsilon}{4}) - \frac{\epsilon}{4} \\ &\stackrel{(57)}{=} P_+(\Omega_+) + P_-(\Omega_-) - \epsilon = \|P\|_{TV} - \epsilon. \end{aligned} \quad (58)$$

◇

A.3 Proof of theorem 2

We start with the following lemma.

Lemma 6. *Let P be a finite signed measure on $(\mathcal{O}, \mathcal{B})$ and $T : \mathcal{O} \rightarrow \mathcal{O}$ a measurable function. Then $P \circ T^{-1}$ is a finite signed measure for which*

$$|P \circ T^{-1}|(B) \leq |P|(T^{-1}B) \quad (59)$$

for all $B \in \mathcal{B}$. In particular,

$$\|P \circ T^{-1}\|_{TV} \leq \|P\|_{TV}. \quad (60)$$

Proof. Note that $P \circ T^{-1} = P_+ \circ T^{-1} - P_- \circ T^{-1}$ is a decomposition into a difference of measures. Because of the uniqueness property of the Jordan decomposition (see th. 4), there is a measure δ such that $P_+ \circ T^{-1} = (P \circ T^{-1})_+ + \delta$ and $P_- \circ T^{-1} = (P \circ T^{-1})_- + \delta$. Therefore $|P \circ T^{-1}|(B) = (P \circ T^{-1})_+(B) + (P \circ T^{-1})_-(B) \leq P_+(T^{-1}B) + P_-(T^{-1}B) = |P|(T^{-1}B)$. $B = \mathcal{O}$ yields the last assertion, as $T^{-1}\mathcal{O} = \mathcal{O}$. $\diamond$

Proof of Th. 2. We recall that, in th. 2, T was supposed to be the shift operator, which is measurable. We observe that

$$\mu P := (P \circ T^{-1}) \quad (61)$$

establishes a linear operator on the vector space of finite, signed measures. Due to lemma 6, (60), it holds that $\|\mu P\|_{TV} \leq \|P\|_{TV}$ for all finite signed measures P , which establishes

$$\|\mu\| \leq 1 \quad (62)$$

where $\|\cdot\|$ is the operator norm associated with the norm of total variation.

Now consider the subspace $\mathcal{P}_P$ of the finite signed measures spanned by all $P \circ T^{-i}$, $i \in \mathbb{N}$ for a given finite signed measure. Note that an equivalent description of finite evolution dimension is just

$$\dim \mathcal{P}_P < \infty. \quad (63)$$

Note further that

$$\mu(\mathcal{P}_P) \subset \mathcal{P}_P. \quad (64)$$

The elementary, linear algebraic lemma 3.2 in [6] states that, given an endomorphism $F : V \rightarrow V$ on a finite-dimensional real- or complex-valued vector space V with $\|F\| \leq 1$, for all $x \in V$ there is an F -invariant $\bar{x} \in V$ such that

$$\lim_{n \rightarrow \infty} \left\| \frac{1}{n} \sum_{k=0}^{n-1} F^k x - \bar{x} \right\| = 0. \quad (65)$$

As all norms are equivalent on V , this applies for arbitrary choices of norms $\|\cdot\|$. Replacing V by $\mathcal{P}_P$, $\|\cdot\|$ by $\|\cdot\|_{TV}$, F by μ and x by P concludes the proof of theorem 2. $\diamond$

B Proof of lemma 7

Lemma 7. *Let P be a discrete random source. Then it holds that*

$$\lim_{t \rightarrow \infty} (H^t(P) - H^t(P \circ T^{-k})) = 0. \quad (66)$$

Proof. Using the notation

$$I_t^k(P) := \frac{1}{t} \sum_{v \in \Sigma^k} \sum_{w \in \Sigma^t} P(vw) \log \frac{P T^{-k}(w)}{P(vw)} \quad (67)$$

and

$$J_t^k(P) := \frac{1}{t} \sum_{v \in \Sigma^k} \sum_{w \in \Sigma^t} P(vw) \log \frac{P(v)}{P(vw)} \quad (68)$$

one obtains

$$H^t(P) + J_t^k(P) \stackrel{(*)}{=} H^{k+t}(P) = I_n^k(P) + H^t(P \circ T^{-k}) \quad (69)$$

where $(*)$ follows from a well known and elementary theorem (e.g. [12], p.22, theorem 2.1) and the second equation is obvious. Because of

$$0 \leq J_t^k(P) \leq \frac{k}{t} H^k(P \circ T^{-t}) S \leq \frac{1}{t} \log |\Sigma^k| \xrightarrow[t \rightarrow \infty]{} 0 \quad (70)$$

and

$$0 \leq I_t^k(X) \leq \frac{1}{t} H^k(X) \leq \frac{1}{t} \log |\Sigma^k| \xrightarrow[t \rightarrow \infty]{} 0, \quad (71)$$

the assertion follows from an application of the sandwich theorem. $\diamond$

Acknowledgment

I would like to thank Ulrich Faigle who considerably contributed to the work presented here. I also would like to thank the unknown reviewers for helpful comments and suggestions.

References

1. D. Aharonov, A. Ambainis, J. Kempe, U. Vazirani, "Quantum walks on graphs", in *Proc. of 33rd ACM STOC, New York, 2001*, pp. 50-59.
2. P. Billingsley, *Ergodic Theory and Information*, Wiley, 1965.
3. L. Breiman, "The individual ergodic theorem of information theory", in *Annals of Mathematical Statistics*, 1957, vol. 28, pp. 809-811.
4. L. Breiman, A correction to 'the individual ergodic theorem of information theory'. *Annals of Mathematical Statistics*, 31:809–810, 1960.

5. U. Faigle, A. Schönhuth, "Quantum predictor models", *Electronic Notes in Discrete Mathematics*, 2006, vol. 25, pp. 149-155.
6. U. Faigle and A. Schoenhuth, "Asymptotic mean stationarity of sources with finite evolution dimension", *IEEE Trans. Inf. Theory*, 2007, vol. 53(7), pp. 2342-2348.
7. H. Federer, *Geometric Measure Theory*. Springer, 1969.
8. R. Gray and L. Davisson, "The ergodic decomposition of stationary discrete random processes", *IEEE Transactions on Information Theory*, 1974, vol. 20(5), pp. 625-636.
9. R.M. Gray and J.C. Kieffer, "Asymptotically mean stationary measures" *Annals of Probability*, 1980, vol. 8, pp. 962-973.
10. Robert M. Gray, *Entropy and Information Theory*. Springer Verlag, 1990.
11. P.R. Halmos, *Measure Theory*. Van Nostrand, 1964.
12. T.S. Han and K. Kobayashi, *Mathematics of Information and Coding*. American Mathematical Society, 2002.
13. A. Heller "On stochastic processes derived from Markov chains", *Annals of Mathematical Statistics*, vol. 36(4), pp. 1286-1291, 1965
14. H. Ito, S.-I. Amari and K. Kobayashi: "Identifiability of hidden Markov information sources and their minimum degrees of freedom", *IEEE Trans. Inf. Theory*, vol. 38(2), pp. 324-333, 1992.
15. S.D. Jacka and G.O. Roberts, "On strong forms of weak convergence" *Stochastic Processes and Applications*, 1997, vol. 67, pp. 41-53.
16. K. Jacobs, "Die bertragung diskreter Informationen durch periodische und fastperiodische Kanäle", 1959, *Mathematische Annalen*, vol. 137, pp. 125-135.
17. K. Jacobs, "ber die Struktur der mittleren Entropie", *Mathematisches Zentralblatt*, 1962, vol. 78, pp. 33-43.
18. P. Jacquet, G. Seroussi, and W. Szpankowski, "On the entropy of a hidden markov process", in *Proc. Data Compression Conf.*, Snowbird, UT, March 2004, pp. 362-371.
19. J.C. Kieffer and M. Rahe, "Markov channels are asymptotically mean stationary", *SIAM J. Math. Anal.*, 1981, vol. 12(3), pp. 293-305.
20. G. Han and B. Marcus, "Analyticity of entropy rate of hidden Markov chains", *IEEE Trans. Inf. Theory*, 2006, vol. 52(12), pp. 5251-5266.
21. G. Han and B. Marcus, "Derivatives of entropy rate in special families of hidden Markov chains", *IEEE Trans. Inf. Theory*, 2007, vol. 53(7), pp. 2642-2652.
22. G. Han and B. Marcus, "Asymptotics of entropy rate of hidden Markov chains at weak black holes", *Proc. IEEE Int. Symp. Inf. Th.*, 2008, pp. 2629-2633.
23. B. McMillan, "The basic theorems of information theory", *Annals of Mathematical Statistics*, 1953, vol. 24, pp. 196-219.
24. Y. Ephraim, N. Merhav, "Hidden Markov processes", *IEEE Trans. on Information Theory*, vol. 48(6), pp. 1518-1569.
25. E. Ordentlich and T. Weissman, "On the optimality of symbol by symbol filtering and denoising", *IEEE Trans. Inf. Theory*, 2006, 52(1), pp. 19-40.
26. A. Schönhuth, "The ergodic decomposition of asymptotically mean stationary random sources", submitted manuscript, <http://arxiv.org/abs/0804.2487>.
27. A. Schönhuth, "On analytic properties of entropy rate", 2006, technical report, ZAIK, University of Cologne.
28. A. Schönhuth, "A simple and efficient solution of the identifiability problem for hidden Markov sources and quantum random walks", ISITA 2008, to appear, <http://arxiv.org/abs/0808.2833>.
29. C. Shannon, "A mathematical theory of communication", *Bell System Technical Journal*, 1948.
30. O. Zuk, E. Domany, I. Kanter, and M. Aizenman, "From finite system entropy to entropy rate for a hidden markov process", *IEEE Signal Processing Letters*, 2006, vol. 13(9), pp. 517-520.

31. O. Zuk, I. Kanter, and E. Domany. The entropy of a binary hidden markov process. *Journal of Statistical Physics*, 2005, vol. 121(3-4), pp. 343-360.